

A table of derivatives and anti-derivatives

This example is based upon a nice example in the Pythontex gallery, see <https://github.com/gpoore/pythontex/>. It uses a tagged block to capture the Sympy output for later use in the body of the LaTeX table.

```
1  from sympy import *
2
3  var('x')
4
5  # Create a list of functions to include in the table
6  funcs = [['sin(x)',r'\sin(x)'], ['cos(x)',r'\cos(x)'], ['tan(x)',r'\tan(x)'],
7            ['asin(x)',r'\arcsin(x)'], ['acos(x)',r'\arccos(x)'], ['atan(x)',r'\arctan(x)'],
8            ['sinh(x)',r'\sinh(x)'], ['cosh(x)',r'\cosh(x)'], ['tanh(x)',r'\tanh(x)']]
9
10 # pyBeg (CalculusTable)
11 for func, eol in funcs:
12     myddx = 'Derivative(' + func + ', x)'
13     myint = 'Integral(' + func + ', x)'
14     print(latex(eval(myddx)) + '&=' + latex(eval(myddx + '.doit()')) + r'\quad & \quad')
15     print(latex(eval(myint)) + '&=' + latex(eval(myint + '.doit()')) + eol)
16 # pyEnd (CalculusTable)
```



```
\begin{align*}
\py{CalculusTable}
\end{align*}
```

$$\frac{d}{dx} \sin(x) = \cos(x)$$

$$\frac{d}{dx} \cos(x) = -\sin(x)$$

$$\frac{d}{dx} \tan(x) = \tan^2(x) + 1$$

$$\frac{d}{dx} \arcsin(x) = \frac{1}{\sqrt{-x^2 + 1}}$$

$$\frac{d}{dx} \arccos(x) = -\frac{1}{\sqrt{-x^2 + 1}}$$

$$\frac{d}{dx} \arctan(x) = \frac{1}{x^2 + 1}$$

$$\frac{d}{dx} \sinh(x) = \cosh(x)$$

$$\frac{d}{dx} \cosh(x) = \sinh(x)$$

$$\frac{d}{dx} \tanh(x) = -\tanh^2(x) + 1$$

$$\int \sin(x) dx = -\cos(x)$$

$$\int \cos(x) dx = \sin(x)$$

$$\int \tan(x) dx = -\log(\cos(x))$$

$$\int \arcsin(x) dx = x \arcsin(x) + \sqrt{-x^2 + 1}$$

$$\int \arccos(x) dx = x \arccos(x) - \sqrt{-x^2 + 1}$$

$$\int \arctan(x) dx = x \arctan(x) - \frac{\log(x^2 + 1)}{2}$$

$$\int \sinh(x) dx = \cosh(x)$$

$$\int \cosh(x) dx = \sinh(x)$$

$$\int \tanh(x) dx = x - \log(\tanh(x) + 1)$$